




Freeform Search

Database: US Pre-Grant Publication Full-Text Database
US Patents Full-Text Database
US OCR Full-Text Database
EPO Abstracts Database
JPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Term: L6 with l3   

Display: 10 **Documents in Display Format:** - **Starting with Number** 1

Generate: ☐ Hit List ☒ Hit Count ☐ Side by Side ☐ Image

Search
Clear
Interrupt

Search History

DATE: Thursday, January 22, 2004 [Printable Copy](#) [Create Case](#)

Set Name Query

side by side

Hit Count Set Name

result set

DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ

<u>L7</u>	L6 with l3	24	<u>L7</u>
<u>L6</u>	plasmid or gene therapy or nucleic or DNA	229527	<u>L6</u>
<u>L5</u>	L4 with l3	8	<u>L5</u>
<u>L4</u>	fusogenic or conjugated or complexed or covalently	188336	<u>L4</u>
<u>L3</u>	L2 with l1	242	<u>L3</u>
<u>L2</u>	cationic or polycationic	156122	<u>L2</u>
<u>L1</u>	polyhistidine or poly-L-histidine or histidine	41549	<u>L1</u>

END OF SEARCH HISTORY

[First Hit](#) [Fwd Refs](#)☐ [Generate Collection](#) [Print](#)

L7: Entry 19 of 24

File: USPT

Apr 18, 2000

DOCUMENT-IDENTIFIER: US 6051429 A

**** See image for Certificate of Correction ****

TITLE: Peptide-enhanced cationic lipid transfections

CLAIMS:

52. The method of claim 51 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, or histidine and where u is an integer from 1 to about 20.

56. The method of claim 55 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, histidine, glycine or proline and where u is an integer from 1 to about 20.

66. The method of claim 62 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, or histidine and where u is an integer from 1 to about 20.

70. The method of claim 69 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, histidine, glycine or proline and where u is an integer from 1 to about 20.

74. The method of claim 73 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, or histidine and where u is an integer from 1 to about 20.

89. The method of claim 88 wherein said nucleic acid-binding group comprises the cationic peptide sequence (Uaa).sub.u where each amino acid, Uaa, in the sequence, independently of any other amino acid in the sequence can be lysine, arginine, ornithine, homoarginine, or histidine and where u is an integer from 1 to about 20.

First Hit

Generate Collection

L7: Entry 14 of 24

File: PGPB

Jul 5, 2001

PGPUB-DOCUMENT-NUMBER: 20010006817
PGPUB-FILING-TYPE: new-utility
DOCUMENT-IDENTIFIER: US 20010006817 A1

TITLE: CELL DELIVERY COMPOSITIONS

PUBLICATION-DATE: July 5, 2001

US-CL-CURRENT: 435/440; 435/325, 435/455, 435/456, 435/458, 435/6, 435/69.1,
435/91.1, 514/44, 530/300, 530/350, 536/23.1

APPL-NO: 09/ 251783 [PALM]
DATE FILED: February 17, 1999
CONTINUED PROSECUTION APPLICATION: CPA

RELATED-US-APPL-DATA:

Application is a non-provisional-of-provisional application 60/075272, filed
February 19, 1998,

PRIORITY INFORMATION

[0001] This application claims priority to the co-pending provisional application
No. 60/075,272 entitled "Cell Delivery Compositions" filed on Feb. 19, 1998, which
is incorporated in its entirety by reference.

[First Hit](#) [Fwd Refs](#)

Generate Collection

Print

L7: Entry 16 of 24

File: USPT

Oct 22, 2002

DOCUMENT-IDENTIFIER: US 6468981 B1

**** See image for Certificate of Correction ****

TITLE: Compositions and methods for targeting pharmaceutically active materials to cells containing androgen receptors

Brief Summary Text (21):

Polycationic salts useful for completing with nucleic acids include salts of cationic polyamines such polylysines, specifically poly-L-lysines, polyarginines, specifically poly-L-arginine, polyhistidine, and protamines.

[First Hit](#) [Fwd Refs](#)☐ [Generate Collection](#)

L7: Entry 16 of 24

File: USPT

Oct 22, 2002

DOCUMENT-IDENTIFIER: US 6468981 B1

**** See image for Certificate of Correction ****

TITLE: Compositions and methods for targeting pharmaceutically active materials to cells containing androgen receptors

Brief Summary Text (21):

Polycationic salts useful for completing with nucleic acids include salts of cationic polyamines such polylysines, specifically poly-L-lysines, polyarginines, specifically poly-L-arginine, polyhistidine, and protamines.

[First Hit](#)

Generate Collection

Print

L7: Entry 14 of 24

File: PGPB

Jul 5, 2001

PGPUB-DOCUMENT-NUMBER: 20010006817
PGPUB-FILING-TYPE: new-utility
DOCUMENT-IDENTIFIER: US 20010006817 A1

TITLE: CELL DELIVERY COMPOSITIONS

PUBLICATION-DATE: July 5, 2001

US-CL-CURRENT: 435/440; 435/325, 435/455, 435/456, 435/458, 435/6, 435/69.1,
435/91.1, 514/44, 530/300, 530/350, 536/23.1

APPL-NO: 09/ 251783 [PALM]
DATE FILED: February 17, 1999
CONTINUED PROSECUTION APPLICATION: CPA

RELATED-US-APPL-DATA:

Application is a non-provisional-of-provisional application 60/075272, filed
February 19, 1998,

PRIORITY INFORMATION

[0001] This application claims priority to the co-pending provisional application
No. 60/075,272 entitled "Cell Delivery Compositions" filed on Feb. 19, 1998, which
is incorporated in its entirety by reference.

First Hit

L7: Entry 14 of 24

File: PGPB

Jul 5, 2001

DOCUMENT-IDENTIFIER: US 20010006817 A1

TITLE: CELL DELIVERY COMPOSITIONS

Detail Description Paragraph:

[0040] Those of ordinary skill in the art will, using known techniques, be able to prepare any of a variety of polyhistidine/polylysine compositions that can readily be tested according to the teachings herein to identify those with desirable delivery characteristics. The compositions must have sufficient polyhistidine composition (including available proton acceptor sites and/or polycationic character) to lyse endosomes, and sufficient polylysine composition to bind to nucleic acids, and condense them if necessary. Thus, the inventive polyhistidine/polylysine composition may comprise any combination of polylysine with polyhistidine, polylysine with histidine, or lysine with polyhistidine, associated with one another covalently or otherwise, so long as the combination is biocompatible and has the endosomolytic and nucleic acid binding/packaging capabilities described herein. As one of ordinary skill in the art will realize, the entire composition (including the bound nucleic acid) must be small enough to be taken up into cells. As mentioned above, endosomal compartments can usually accept entities up to about 150 nm in size.

Detail Description Paragraph:

[0103] The ability of the packaging agent to bind DNA can be assessed by monitoring complex formation with DNA using gel electrophoresis. The mobility of DNA on the gel will be retarded by complex formation, and the absence of any mobility of DNA on the gel suggests the complexation of all of the DNA. Preferably, complexation of DNA and the cationic polymer occurs as a ratio of 1:1 DNA/cationic polymer, and most preferably at a ratio of 1:3 DNA/cationic polymer as shown in FIG. 13 and 14 for DNA transferrin-polylysine and DNA/G-pHis mixtures. FIG. 15 depicts the gel electrophoresis of DNA/p-His mixtures and shows complexation at a weight:weight ratio of 1:0.5 DNA/p-His. Condensing of plasmid DNA can also be monitored by observing the ethidium bromide exclusion. For example, if gluconylated polyhistidine is used as the cationic polymer, the gluconylated polyhistidine efficiently condenses DNA at pH 5 where the gluconylated polyhistidine is .about.45% protonated. DNA is not condensed as effectively, however, at pH 7.4 where gluconylated polyhistidine is .about.5% protonated, as shown in FIG. 11.



STIC Search Report

Biotech-Chem Library

STIC Database Tracking Number: 112371

TO: Dave Nguyen

Location: rem2d31

Art Unit: 1623

Jan 22, 2004

Case Serial Number: 10/018103

From: P. Sheppard

Location: Remsen Building

Phone: (571) 272-2529

sheppard@uspto.gov

Search Notes

112371

STIC-Biotech/ChemLib

From: Page, Thurman
Sent: Saturday, January 17, 2004 11:39 AM
To: STIC-Biotech/ChemLib
Cc: Nguyen, Dave; Page, Thurman
Subject: FW: Rush Search request 10/018,103
Importance: High

RUSH SEARCH APPROVED

Thurman K. Page
SPE Art Units 1615 & 1616
Technology Center 1600

-----Original Message-----

From: Nguyen, Dave
Sent: Friday, January 16, 2004 9:17 PM
To: Page, Thurman
Cc: STIC-Biotech/ChemLib
Subject: Rush Search request 10/018,103

This case is due this bi-week. Please rush. Please do a polypeptide/peptide search on SEQ ID NOS: 4-6.

Thanks,
Dave Nguyen
Art Unit: 1632
Ramsen Building
2D31
571-272-0731

Searcher: Sheppard
Phone: _____
Location: _____
Date Picked Up: _____
Date Completed: 1/22/04
Searcher Prep/Review: _____
Clerical: _____
Online time: _____

TYPE OF SEARCH:
NA Sequences: _____
AA Sequences: _____
Structures: _____
Bibliographic: _____
Litigation: _____
Full text: _____
Patent Family: _____
Other: _____

VENDOR/COST (where applic.)
STN: _____
DIALOG: _____
Questel/Orbit: _____
DRLink: _____
Lexis/Nexis: _____
Sequence Sys.: _____
WWW/Internet: _____
Other (specify): _____